
### **H4** - Preparation for farrowing



A farrowing pen, ready for farrowing to take place

## A good farrowing process requires thorough preparation:

- 1. Transfer the sow to the farrowing pen min. 1 week before farrowing.
- 2. Supply 2.8-3.0 FUsow/day the day before expected farrowing.
- 3. Remove all dirt behind the sow the last 2 days before expected farrowing until farrowing is complete.
- 4. Flush the piglets' water supply before farrowing starts. For instance, put a clothes-peg on the drinker for a few minutes.
- 5. Supply extra water to the sow if required.
- 6. Litter the creep area before farrowing.
- 7. Adjust the farrowing crate inwards.
- 8. Supply nesting material once a day a week before farrowing.
- 9. Install a heat lamp and turn it on the day before farrowing.
- Prepare boards for the creep areas, equipment for obstetric aid and medication.
- 11. Act calmly around sows about to farrow.



The heat lamp is missing and the board in front of the creep area is up

# Inadequate preparation may result in: Incorrect feeding

 Restless and stressed sows, and prolonged farrowings.

#### Poor environment for the piglets

 Cold/supercool piglets are weakened and have high mortality rates

#### Poor hygiene

- Increased risk of the piglets becoming sick.
- · Weak piglets.

#### Insufficient time for obstetric aid

- More stillborn piglets
- Weakened sows
- Piglets born weak.